

Tsemo Aristide  
 103-135 Leeward Glenway  
 North York  
 M3C 2Z6  
 tsemoaristide@hotmail.com

## KAHLER, POISSON GEOMETRY OF CR LIE GROUPS.

*"Las frutas de la honestidad se recogen  
 en muy poco tiempo y duran para siempre."  
 Colombian Proverb.*

### Abstract.

A Cauchy Riemann (CR) Lie group is a Lie group  $G$  which Lie algebra  $\mathcal{G}$  has a vector subspace  $\mathcal{H}$  endowed with an endomorphism  $j$  such that  $j^2 = -Id$ , and for each elements  $x, y$  in  $\mathcal{H}$ , we have  $[j(x), j(y)] - [x, y]$  is an element of  $\mathcal{H}$ , and  $[j(x), j(y)] - [x, y] = j([j(x), y] + [x, j(y)])$ . In this paper we study the geometry of a CR Lie groups  $G$  when its Lie algebra  $\mathcal{G}$  is endowed with more geometric structures compatible with  $j$ , as kahler, and poisson type structures.

### 0. Introduction.

Let  $(M, j)$  be a complex manifold, and  $H$  an hypersurface of  $M$ , for each element  $x$  of  $H$  the tangent space  $TH_x$  of  $H$  at  $x$  is endowed with a maximal complex vector space  $E_x = TH_x \cap j(TH_x)$ . The collection of vector spaces  $E_x$  defines a vector bundle  $E$  such that for sections  $X$ , and  $Y$  of  $E$ , we have:

$$[j(X), j(Y)] - [X, Y] \in E$$

since  $(M, j)$  is a complex structure, we also have:

$$[j(X), j(Y)] - [X, Y] = j([j(X), Y] + [X, j(Y)]).$$

More generally, a Cauchy Riemann (CR) structure on a manifold  $M$ , is defined by a subbundle  $H$  of  $TM$  endowed with an endomorphism  $j$  such that:

$$j^2 = -Id_H$$

For each sections  $X$  and  $Y$  of  $H$ , we have:

$$[j(X), j(Y)] - [X, Y] \in H$$

$$[j(X), j(Y)] - [X, Y] = j([j(X), Y] + [X, j(Y)]),$$

The notion of CR manifolds is studied by many authors see[3]

In this paper we study Left-invariant  $CR$ -structures on Lie groups compatible with geometric properties as poisson, and kahler type properties. More precisely:

**Definition 0.1.**

A kahlerian complex real Lie group  $(G, \mathcal{H}, j, <, >)$ , is a Lie group  $G$  endowed with a CR structure defined by the vector subspace  $\mathcal{H}$ ,  $j$  is an endomorphism of  $\mathcal{G}$  which image is  $\mathcal{H}$ . We suppose that the following properties are verified:

1.  $j$  preserves  $\mathcal{H}$ , the restriction of  $j^2$  to  $\mathcal{H}$  is  $-Id$ .
2.  $[X, Y] - [jX, jY] \in \mathcal{H}$  if  $X, Y \in \mathcal{H}$
3.  $[j(X), j(Y)] = [X, Y] + j([X, j(Y)] + [j(X), Y])$  if  $X, Y \in \mathcal{H}$

Moreover we will suppose that there exists a left-invariant riemannian metric on  $G$ , defined by a scalar product on  $\mathcal{G}$  such that  $\omega = <, j >$  is closed. This means that  $\omega$  is antisymmetric and

$$\omega([x, y], z) + \omega([z, x], y) + \omega([y, z], x) = 0$$

We remark that the restriction of  $\omega$  to  $\mathcal{H}$  is not degenerated.

We show results related to those structures, for example, we show that a semi-simple kahlerian CR Lie group such that the codimension of  $\mathcal{H}$  is 1 is locally isomorphic to  $so(3)$  or  $sl(2)$ .

**1. The structure of kahlerian Lie groups.**

Let  $G$  be Lie group. A left symmetric structure on  $G$ , is defined by a product on its Lie algebra  $(\mathcal{G}, [,])$

$$\mathcal{G} \times \mathcal{G} \longrightarrow \mathcal{G}$$

$$(x, y) \longrightarrow xy$$

which verified

$$xy - yx = [x, y]$$

and

$$x(yz) - (xy)z = y(xz) - (yx)z.$$

This is equivalent to endows  $G$  with a left invariant connection which curvature and torsion forms vanish identically.

In this part we consider a CR kahlerian Lie group  $(G, \mathcal{H}, j, <, >)$ . The Lie algebra  $\mathcal{G}$  of  $G$ , will be call a CR kahlerian Lie algebra.

We will define the following product on  $\mathcal{H}$ :

For  $x, y, z \in \mathcal{H}$ , we set

$$\omega(xy, z) = -\omega(y, [x, z]) = \omega([x, z], y) = < [x, z], j(y) >$$

**Proposition 1.1.**

For every  $x, y, u$  and  $z$  in  $\mathcal{H}$  we have

$$(1) \quad \omega(xy - yx, u) = \omega([x, y], u)$$

and if the bracket  $[x, y]' = xy - yx$  satisfies the jacobi identity

$$(2) \quad x(yz) - (xy)z = y(xz) - (yx)z$$

**Proof.**

The proof almost copy the one of left-invariant symplectic structures on Lie groups.

Let  $u$  be an element of  $\mathcal{H}$ , for  $x, y, z$  in  $\mathcal{H}$ , we have  $\omega(xy - yx, u) = -\omega(y, [x, u]) + \omega(x, [y, u])$  then (1) follows from the definition of  $\omega$  (the closed property).

Now we prove the second assertion:

$$\omega(x(yz), u) = -\omega(yz, [x, u]) = \omega(z, [y, [x, u]]')$$

We also have:

$$\omega((xy)z, u) = -\omega(z, [xy, u]) = \omega(z, [u, xy]) = -\omega(uz, xy)$$

This implies that

$$\omega(x(yz) - (xy)z, u) - \omega(y(xz) - (yx)z, u) =$$

$$\omega(z, [y, [x, u]] - [x, [y, u]]) + \omega(uz, xy - yx)$$

The property (1) implies that:

$$\omega(uz, xy - yx) = \omega(uz, [x, y]') = -\omega(z, [u, [x, y]]')$$

We deduce that

$$\omega(x(yz) - (xy)z, u) - \omega(y(xz) - (yx)z, u) = \omega(z, [y, [x, u]]' + [x, [u, y]]' + [u, [y, x]]') = 0$$

**Corollary 1.2.**

Let  $\mathcal{G}, \mathcal{H}, j$  be a kahlerian cr-algebra, then the product  $(x, y) \rightarrow xy - yx = [x, y]'$  defined on  $\mathcal{H}$  a structure of a kahlerian Lie algebra if it satisfies the Jacobi identity.

**Proof.**

We deduce from the property (2) that the product defined on  $\mathcal{H}$ ,  $(x, y) \rightarrow xy$  endows  $\mathcal{H}$  with a structure of Left symmetric algebra which underlying Lie algebra is  $[\cdot, \cdot]'$ , the morphism  $j$  defines also a complex structure on  $H$ , and the scalar product  $\langle, \rangle$  a kahlerian structure.

**Proposition 1.3.**

Consider the vector subspace  $\mathcal{L}$  such that for each  $x \in \mathcal{L}$ , we have  $\omega(x, \mathcal{G}) = 0$ ,  $\mathcal{L}$  is a Lie subalgebra of  $\mathcal{G}$  and is the  $\langle, \rangle$  orthogonal vector space of  $\mathcal{H}$ .

**Proof.**

Let  $x$  and  $y$  two elements of  $\mathcal{L}$ , for every element  $z$  of  $\mathcal{G}$  we have:

$$\omega([x, y], z) = \omega(y, [z, x]) + \omega(x, [y, z])$$

since  $x$  and  $y$  are elements of  $\mathcal{L}$ , we deduce that  $\omega([x, y], z) = 0$ , and that  $[x, y] \in \mathcal{L}$ .

For  $x \in \mathcal{L}$ , and  $y \in \mathcal{H}$ , we have:

$$\omega(x, y) = \langle x, j(y) \rangle = 0,$$

Since the restriction of  $j$  to  $\mathcal{H}$  is an automorphism, we deduce that  $\mathcal{L}$  and  $\mathcal{H}$  are  $\langle, \rangle$  orthogonal each other

**Proposition 1.4.**

Let  $L$  be the Lie subgroup which Lie algebra is  $\mathcal{L}$ , and  $M$  the right quotient  $G/L$ , then  $M$  is a kahlerian manifold.

**Proof.**

The fact that  $\mathcal{H}$  and  $\mathcal{L}$  are orthogonal each other implies that the riemannian metric  $\langle, \rangle$  gives rise to a metric  $\langle, \rangle'$  of  $M$ , the morphism  $j$  also gives rise to a complex structure  $j'$  of  $M$ . Denote by  $p$  the projection  $p : G \rightarrow G/L$ , we have  $\omega = p^* \langle, j' \rangle'$ . This implies that  $\omega' = \langle, j' \rangle'$  is a symplectic form defined on  $M$ , thus  $(M, \omega', j')$  is a kahlerian manifold.

Let  $(\mathcal{H}, [, j', \langle, \rangle')$  be a kahlerian Lie algebra, and  $V$  a vector space. Supposed defined a Lie algebra structure on  $\mathcal{G} = \mathcal{H} + V$  such that there exists a map:

$$\alpha : \mathcal{H} \times \mathcal{H} \rightarrow V$$

such that for  $x, y \in \mathcal{H}$  we have  $[x, y] = [x, y]' + \alpha(x, y)$ , suppose that  $\alpha(j(x), j(y)) = \alpha(x, y)$ .

We suppose also that there exists a scalar product  $\langle, \rangle$  on  $\mathcal{G}$  which extends  $\langle, \rangle'$  such that  $\mathcal{H}$  and  $V$  are orthogonal, we also extend  $j'$  to an endomorphism  $j$  of  $\mathcal{G}$  such that  $j(V) = 0$ . We suppose that the form  $\omega = \langle, j \rangle$  is closed, then  $(\mathcal{G}, j, \langle, \rangle)$  is a  $CR$ -kahlerian algebra.

Remark that the fact that  $\mathcal{G}$  is a kahlerian  $CR$ -Lie algebra implies the following property:

$$\oint \alpha([x, y]', z) + [\alpha(x, y), z] = 0$$

Remark that if  $\mathcal{H}$  is a sub Lie algebra of  $\mathcal{G}$ , then its symplectic structure defined on  $\mathcal{G}$  a left invariant Poisson structure.

**Examples of kahlerian  $CR$ -structures.**

1.

Consider the  $n$ -dimensional commutative Lie algebra  $\mathbb{R}^n$  endowed with its flat riemannian metric  $\langle, \rangle$ , and  $V$  an even dimension subspace of  $\mathbb{R}^n$  endowed with a linear map  $j$  such that  $j^2 = -id$ , then  $(\mathbb{R}^n, V, j, \langle, \rangle)$  is a Lie kahlerian  $CR$ -algebra.

2.

Consider the semi-simple algebra  $so(3)$ , and  $(e_1, e_2, e_3)$  its basis in which its Lie structure is defined by

$$[e_1, e_2] = e_3, [e_1, e_3] = -e_2, [e_2, e_3] = e_1$$

On  $Vect(e_1, e_2)$  the subspace generated by  $e_1$  and  $e_2$ , we consider the linear map  $j$  defines by  $j(e_1) = e_2$  and  $j(e_2) = -e_1$ .

Let  $\langle, \rangle$  be the scalar product defined on  $so(3)$  by  $\langle e_i, e_j \rangle = \delta_{ij}$ ,

The family  $(so(3), V, j, \langle, \rangle)$  is a  $cr$ -kahlerian Lie algebra.

**Proposition 1.6.**

Let  $Z(\mathcal{G})$  be the center of  $\mathcal{G}$ , then  $U = (Z(\mathcal{G}) \cap \mathcal{H}) + j(Z(\mathcal{G}) \cap \mathcal{H})$  is a Lie commutative algebra and for every element  $z$  of  $U$ ,  $adz(\mathcal{H}) \subset \mathcal{H}$ .

**Proof.**

Let  $z$  and  $z'$  be elements of  $Z(\mathcal{G}) \cap \mathcal{H}$ , we have  $[j(z), j(z')] = [z, z'] + j([z, j(z')]) + [j(z), z']$ . Since  $z$  and  $z'$  are in the center of  $\mathcal{G}$ , we deduce that  $[j(z), j(z')] = 0$ .

The fact that  $[z, \mathcal{H}] \subset \mathcal{H}$  follows from the fact that for  $x, y$  in  $\mathcal{H}$ , we have  $[x, j(y)] + [j(x), y]$  is an element of  $\mathcal{H}$ .

**Proposition 1.7.**

Let  $\mathcal{G}$  be a  $CR$ -algebra, suppose that there exists an ideal  $I$  supplementary to  $\mathcal{H}$ , then  $\mathcal{H}$  is endowed with a complex Lie structure.

**Proof.**

The projection  $p : \mathcal{G} \rightarrow \mathcal{H}$  parallel to  $I$  defines on  $\mathcal{H}$  a Lie-complex structure.

Conversely suppose given an extension

$$0 \longrightarrow I \longrightarrow \mathcal{G} \longrightarrow \mathcal{U} \longrightarrow 0$$

where  $\mathcal{U}$  is a Lie algebra endowed with a complex structure, then a supplementary space  $\mathcal{H}$  of  $I$  defines a  $CR$ -structure on  $\mathcal{G}$  if and only if for every  $x, y$  in  $\mathcal{H}$ ,  $[jx, jy] - [x, y]$  is an element of  $\mathcal{H}$  and  $[j(x), j(y)] - [x, y] = j([j(x), y] + [x, j(y)])$ , where  $j$  is the pulls-back of the complex structure of  $\mathcal{U}$ .

**Proposition 1.8.**

Let  $(G, \mathcal{H}, \langle, \rangle, j)$  be a  $CR$  kahlerian Lie group. Suppose that  $\mathcal{H}^\perp = \ker j$  is an ideal, then  $\mathcal{G}$  is not semi-simple.

**Proof.**

We have seen that if  $\mathcal{H}^\perp$  then  $\mathcal{H}$  is endowed with a structure of a kahlerian Lie algebra which is known not be semi-simple. Since the quotient of  $\mathcal{G}$  by  $\mathcal{H}^\perp$  is isomorphic as a Lie algebra to  $\mathcal{H}$ , we deduce that  $\mathcal{G}$  is not semi-simple.

**Proposition 1.9.**

Suppose that  $\mathcal{H}^\perp$  is an ideal, then  $G$  is the product of a family of groups  $G_i$ , where  $G_0$  is flat, and for  $i \geq 1$ , the holonomy of the riemannian structure of  $G_i$  is irreducible. Each group  $G_i$  is endowed with a kahlerian CR-structure defined by the subspace  $\mathcal{H}_i$  of the Lie algebra  $\mathcal{G}_i$  of  $G_i$  such that the sum of the dimension of  $\mathcal{H}_i$  is  $\mathcal{H}$ .

**Proof.**

Suppose that  $\mathcal{H}^\perp$  is an ideal, then the quotient  $\mathcal{L}$  of  $\mathcal{G}$  by  $\mathcal{H}^\perp$  is a kahlerian Lie algebra. The theorem (Lichnerowicz Medina [12]), implies that  $\mathcal{L} = \sum \mathcal{L}_i$  where each  $\mathcal{L}_i$  is a kahlerian Lie algebra. Now consider the De Rham decomposition of  $G$  as a product of groups  $G_i$ . The projection  $G \rightarrow L$  respect this decomposition. Suppose that the projection  $p : \mathcal{G}_i \rightarrow \mathcal{L}_i$  is not trivial, then the orthogonal of the kernel of  $p$  defines  $\mathcal{H}_i$ .

Consider the set of functions  $C^\infty(G_H)$  defined on  $G$  such that  $d^n f \in S^n(T^*\mathcal{H})$  where  $T\mathcal{H}$  is the left invariant distribution on  $G$  defined by  $\mathcal{H}$ . For each  $f \in C^\infty(G_H)$ , there exists a vector field  $X_f$  such that

$$\omega(X_f, \cdot) = df$$

we will denote by  $\{f, g\} = \omega(X_f, X_g)$ .

**Proposition 1.10.**

The algebra  $(C^\infty(G_H), \{, \})$  is a Poisson algebra, i.e  $\{, \}$  verifies the Jacobi identity.

## 2. Homogeneous Kahler CR manifold.

Let  $(G, \mathcal{H}, j, <, >)$  be a kahler CR Lie group and  $\Gamma$  a cocompact discrete subgroup of  $G$ , the manifold  $M = G/\Gamma$  inherits a CR structure from  $G$ .

The orbits of the left action of the group  $L$  on  $G$  defines a foliation  $\mathcal{F}_L$  on  $M$ . (The Lie algebra of  $L$  is  $\mathcal{H}^\perp$ , orthogonal).

**Proposition 2.1.**

The orbits of the foliation  $\mathcal{F}_L$  are closed if and only if the group  $\Gamma L$  is closed in  $G$ , in this case  $M$  is the total space of a fibration over a kahler manifold.

**Proof.**

Suppose that the group  $\Gamma L$  is closed, then the quotient  $G/\Gamma L$  is a Kahlerian manifold  $N$ , the projection map  $M \rightarrow N$  induced by the identity map of  $G$  which is the given fibration.

Conversely suppose that the orbits of the foliation  $\mathcal{F}_L$  are closed. Consider a sequence  $g_n$  of  $\Gamma L$  which converges towards the element  $g$  of  $G$ . Consider a neighbourhood  $U$  of  $g$  such that the restriction of the projection  $p : G \rightarrow M$  to  $U$  is injective, then  $p(g)$  is an element of the adherence of  $p((\Gamma L)e) = \mathcal{F}_{p(e)}$ , where  $e$  is the neutral element of  $G$ . Since we have supposed that the leaves of  $\mathcal{F}_L$  are closed,  $p(g)$  is an element of  $\mathcal{F}_{p(e)}$  which means that  $\Gamma L$  is closed in  $G$ . We can thus define the Kahler manifold  $G/\Gamma L$ .

### Deformation of CR kahlerian structures of homogeneous manifolds.

Let  $(G, \mathcal{H}, j, <, >)$  be a Lie group endowed with a CR kahlerian structure. Consider a cocompact subgroup  $\Gamma$  of  $G$  the manifold  $M = G/\Gamma$  inherits from  $G$  a CR structure. In this section, we will define the deformation of those structures from two points of view.

Supposed fixed the CR kahlerian structure of  $G$ , and a compact manifold  $M$ , Let  $\Gamma$  be a group we consider the set of representations  $R(\Gamma, G)$  such that for each  $u \in R(\Gamma, G)$ ,  $u$  is injective and  $G/u(\Gamma)$  is a compact manifold.

To elements  $u$  and  $u'$  of  $R(\Gamma, G)$  will be said equivalent if and only there exists an element  $g$  of  $G$  such that  $u' = gug^{-1}$ . We denote by  $Def_1(\Gamma, G, \mathcal{H}, j, <, >)$  the space of equivalence classes of those CR kahlerian structures.

Now consider  $RCK(G)$ , the set of real complex kahlerian structures of  $G$ , then for a cocompact subgroup  $\Gamma$  of  $G$   $M = G/\Gamma$  inherits a CR kahlerian structure for each element  $u$  of  $RCK(G)$  denotes by  $(M, u)$ . We will say that  $(M, u_1)$  is equivalent to  $(M, u_2)$  if and only if there exists an isomorphism of CR kahlerian complex manifolds between  $(M, u_1)$  and  $(M, u_2)$ . and denote by  $Def_2(M, G)$  the set of those CR structures.

### 3. Kahlerian codimension 1 CR-structures.

#### Theorem 3.1.

*Suppose that the codimension of  $\mathcal{H}$  is  $l$ , and  $G$  is semi-simple then  $G$  is a Lie group of rank  $\leq l$ .*

#### Proof.

Let  $(\mathcal{G}, \mathcal{H}, j, <, >)$  be a Lie semi-simple algebra endowed with a codimension  $l$  CR-kahlerian structure. This means that the codimension of  $\mathcal{H}$  in  $\mathcal{G}$  is  $l$ .

The map  $\mathcal{G} \rightarrow \mathcal{G}^*$ ,

$$X \longrightarrow \omega(X, \cdot)$$

is a 1-cocycle for the coadjoint representation. Since  $\mathcal{G}$  is semi-simple, this cocycle is trivial. There exists an element  $\alpha$  of  $\mathcal{G}^*$  such that for  $x, y \in \mathcal{G}$  we have:

$$\omega(x, y) = \alpha([x, y])$$

Let  $K$  be the Killing form of  $\mathcal{G}$ , there exists  $X \in \mathcal{G}$  such that for each  $Y \in \mathcal{G}$  we have:

$$K(X, Y) = \alpha(Y)$$

The Lie algebra  $\mathcal{L} = \{x, : \omega(x, \mathcal{G}) = 0\}$  is a dimension  $l$  subalgebra of  $\mathcal{G}$  since the codimension of  $\mathcal{H}$  is  $l$ . The Lie algebra  $\mathcal{L}$  is the Lie algebra of the subgroup  $L$  of  $G$  which preserves  $\alpha$ ,  $L$  is also the subgroup which preserves  $X$  since  $K$  is invariant by the adjoint representation. We deduce that the rank of  $G$  is less or equal than  $l$ , and then that  $G$  is isomorphic to  $sl(2)$  or  $so(3)$  if the codimension of  $\mathcal{H}$  is 1.

**Corollary 3.2.**

Suppose that  $\mathcal{G}, \mathcal{H}, <, >, j$  is a semi-simple codimension 1  $cr$ -structure (the codimension of  $\mathcal{H}$  is 1), then  $\mathcal{G}$  is  $so(3)$  or  $sl(2)$ .

**4. Poisson  $CR$ -structures.**

Let  $M$  be a manifold, and  $TH$  and  $TU$  two supplementary subbundles of its tangent bundle  $TM$ .

**Definition 4.1.**

An  $(TH, TU)$ -pseudo-Poisson structure on  $M$  is defined by a bivector  $\Lambda \in \Lambda^2 TM$  such that

$$[\Lambda, \Lambda] \in TU\Lambda^2 TM$$

where  $[\Lambda, \Lambda]$  is the schouten product of  $\Lambda$  by  $\Lambda$ . The bivector  $\Lambda$  defined on  $C^\infty(M)$  the bracket  $\{, \}$  by the formula

$$\{f, g\} = \Lambda(df, dg)$$

A morphism  $f : (M, \Lambda) \rightarrow (M', \Lambda')$  is a differentiable map which commutes with  $\{, \}$ .

Suppose that the distribution  $TH$  defines on  $M$  a  $cr$ -structure, we will say that  $(M, TH, TU, j)$  defines a pseudo-Poisson  $cr$ -structure on  $M$ , if  $j$  preserves  $\Lambda$ .

Let  $G$  be a Lie group which Lie algebra is  $\mathcal{G}$ . Consider a subspace  $H$  of  $\mathcal{G}$  and  $U$  a supplementary space to  $H$ . The vector spaces  $H$  and  $U$  define on  $G$  right invariant distributions  $TH$  and  $TU$ .

**Definition 4.2.**

A pseudo-Lie Poisson structure on  $G$  is a bivector  $\Lambda \in \Lambda^2 TG$  such that

$$[\Lambda, \Lambda] \in TU\Lambda^2 TG.$$

Moreover we suppose that  $\Lambda$  is multiplicative i.e that the product  $G \times G \rightarrow G$  is a morphism of pseudo-Poisson structures.

The bracket  $\{, \}$  defined by  $\Lambda$  satisfies the following properties:

$$\{f, f'\}(xy) = \{f \circ L_x, f' \circ L_x\}(y) + \{f \circ R_y, f' \circ R_y\}(x).$$

If we denote by  $T_u L_x$  the differential of  $L_x$  in  $u$ , we have

$$\Lambda(xy) = T_y L_x \Lambda_y + T_x R_y \Lambda_x$$

Consider the tensors  $\Lambda_R(x) = T_x R_{x^{-1}} \Lambda_x$  and  $\Lambda_L(x) = T_x L_{x^{-1}} \Lambda_x$

**Proposition 4.3.**

The fact that  $\pi$  is multiplicative is equivalent to

1.  $\pi_R$  is a 1-cocycle for the adjoint representation  $G \rightarrow \Lambda^2 \mathcal{G}$ , i.e  $\pi_R(xy) = \pi_R(x) + Ad_x(\pi_R(y))$ .



2.  $\pi_L$  is a 1-cocycle for the adjoint action of the opposite group of  $G$  in  $\Lambda^2 G$  i.e  $\pi_L(xy) = \pi_L(y) + Ad_{y^{-1}}(\pi_L(x))$ .

Let  $r$  be an element of  $\Lambda^2 \mathcal{G}$ ,  $r_-$ , and  $r_+$  the left and right invariant tensors defined by  $r$ . We denote by  $\pi$  the tensor  $r_+ - r_-$ .

The tensor  $\pi$  defines a pseudo-Poisson structure if and only if  $[\pi, \pi] = [r, r]_+ - [r, r]_- \in TU\Lambda^2 TG$ , or equivalently if

$$Ad_x[r, r] - [r, r] \in U\Lambda^2 \mathcal{G}.$$

Suppose that  $H$  defines on  $G$  a  $cr$ -structure. We will say that  $(G, H, \Lambda, j)$  is a  $cr$ -Poisson structure if  $j$  preserves  $\Lambda$ .

Moreover we assume that  $\Lambda$  is invariant by  $j$ .

**Remark.**

Suppose defined the  $cr$ -Poisson structure  $(\mathcal{G}_1, \mathcal{H}_1, j_1, \Lambda_1)$  and  $(\mathcal{G}_2, \mathcal{H}_2, j_2, \Lambda_2)$ , then the tensor  $\Lambda_1 \times \Lambda_2$  defines a  $cr$ -structure  $(\mathcal{G}_1 \times \mathcal{G}_2, \mathcal{H}_1 \times \mathcal{H}_2, \Lambda \times \Lambda_2, j_1 \times j_2)$  called the product of the Poisson  $cr$ -structures.

**References.**

1. Aubert, A. Structures affines et pseudo-metriques invariantes a gauche sur des groupes de Lie.  
Thesis Universite de Montpellier II 1996
2. Benson, C. Gordon C. Kahler and symplectic structures on nilmanifolds.  
Topology 27 (1988) 513-519
3. Bland, J. Epstein, C. Embeddable CR structures and deformation of pseudo-convex surfaces.  
J. Algebraic Geometry 5 (1996) 277-368
4. Boothby, W. Wang, H. On contact manifolds.  
Ann. Math. 68. (1958) 721-734
5. Dardie J-M, Groupes de Lie symplectiques ou kahleriens et double extension.  
Thesis Universite de Montpellier II 1993
6. Diatta, A. Geometrie de Poisson et de Contact des espaces homogenes  
Thesis Universite de Montpellier II 2000
7. Domini, S. Gigante, G. Classification of left invariant CR structures on  $Gl(3, \mathbb{R})$ .  
Rivist. Mat. Univ di Parma.
8. Dorfmeister, J. Nakajima, K. The fundamental conjecture for homogeneous kahler manifolds.  
Acta Mathematica 161 (1986) 189-208
9. Gauthier, D. Lie CR algebras.

- Diff. Geometry and its applications (1997) 365-376
10. Gigante, G. Tomassini. CR structures on a real Lie algebra.  
Adv. Math (1992) 67-81
  11. Lichnerowicz, A. Les groupes Kahleriens,  
Symplectic geometry and mathematical physics (Edit P. Donato et al). Pro.  
Math 99
  12. Lichnerowicz A, Medina, A. On Lie groups with left invariant symplectic  
or kahlerian structures.  
Lett. Math. Phs. 16 (1988)225-235
  13. Milnor, J. Curvature of left invariant metric on Lie groups.  
Adv. Math. (1976) 293-329
  14. Tsemo, A. Kahlerian, Poisson CR structures on Lie groups and the  
method of double extension. In preparation.